

Non-Abelian Generalization of Off-Diagonal Geometric Phases

David Kult^{1*}, Johan Åberg^{2†}, and Erik Sjöqvist^{1‡}

¹*Department of Quantum Chemistry, Uppsala University, Box 518, Se-751 20 Uppsala, Sweden.*

²*Centre for Quantum Computation, Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, United Kingdom.*

(Dated: February 1, 2008)

If a quantum system evolves in a noncyclic fashion the corresponding geometric phase or holonomy may not be fully defined. Off-diagonal geometric phases have been developed to deal with such cases. Here, we generalize these phases to the non-Abelian case, by introducing off-diagonal holonomies that involve evolution of more than one subspace of the underlying Hilbert space. Physical realizations of the off-diagonal holonomies in adiabatic evolution and interferometry are put forward.

PACS numbers: 03.65.Vf

I. INTRODUCTION

A quantal system that fails to return to its initial state after some prescribed elapse of time may acquire a well-defined geometric phase [1]. An interesting feature of this noncyclic geometric phase is that it becomes undefined when the initial and final states are orthogonal. This gives rise to a nodal point structure that can be monitored experimentally in a history-dependent manner [2, 3, 4, 5]. In the hope to recover some of the lost interference information at the nodal points of the noncyclic geometric phase, Manini and Pistolesi [6] introduced off-diagonal geometric phases for adiabatic evolutions of pure states. These quantities may be defined in cases where the standard geometric phase is not. The adiabatic requirement on the evolution in ref. [6] was lifted by Mukunda *et al.* [7], and Hasegawa *et al.* [8, 9] provided an experimental verification of the second order off-diagonal geometric phase for neutron spin. Theories for off-diagonal phases and holonomies for mixed quantal states have been developed [10, 11, 12, 13].

Wilczek and Zee [14] showed that the geometric phase factor generalizes to a unitary state change, often referred to as a non-Abelian quantum holonomy, when considering cyclic adiabatic evolution governed by a degenerate Hamiltonian. The relevance of non-Abelian holonomies for universal fault tolerant quantum computation has been demonstrated in refs. [15, 16, 17, 18, 19, 20]. The non-Abelian quantum holonomies have been generalized to nonadiabatic [21], discrete [22], and noncyclic evolutions [23, 24]. As for the geometric phase, the holonomy may be undefined when the evolution is noncyclic. In the non-Abelian case we also have the additional possibility that the holonomy is partially defined [24].

In this Letter, we extend ref. [6] and introduce non-Abelian off-diagonal holonomies. We demonstrate that the off-diagonal holonomies retain holonomy information

when the standard noncyclic ones [23, 24] are undefined. We also suggest physical realizations of the off-diagonal holonomies using interferometry, both in adiabatic and nonadiabatic settings.

II. OFF-DIAGONAL HOLONOMIES

Consider a smoothly parameterized decomposition

$$\mathcal{H} = \mathcal{H}_1(s) \oplus \cdots \oplus \mathcal{H}_\eta(s), \quad s \in [0, 1], \quad (1)$$

of an N -dimensional Hilbert space $\mathcal{H}$ into η mutually orthogonal subspaces. It is to be noted that this parameter-dependent decomposition can arise in different ways, e.g., it may consist of the instantaneous eigenspaces of some s -dependent Hamiltonian [14], or of some arbitrary decomposition that evolves under the Schrödinger equation [21]. Assume that $\dim[\mathcal{H}_l(s)] = n_l$, $\forall s \in [0, 1]$, and $l = 1, \dots, \eta$. Thus, each family $\mathcal{H}_l(s)$ of subspaces defines a curve $\mathcal{C}_l$ in the Grassmann manifold $\mathcal{G}(N; n_l)$, i.e., the set of n_l -dimensional subspaces in the N -dimensional Hilbert space [25]. For each such curve, we introduce the quantities

$$\Gamma_l = \lim_{\delta s \rightarrow 0} P_l(1)P_l(1 - \delta s) \cdots P_l(\delta s)P_l(0), \quad (2)$$

where $P_l(s)$ is the projection operator onto the subspace $\mathcal{H}_l(s)$. This definition makes Γ_l explicitly gauge invariant. Note also that the limit $\delta s \rightarrow 0$ in Eq. (2) makes Γ_l uniquely determined for any sufficiently smooth curve $\mathcal{C}_l$ in the Grassman manifold. We furthermore define

$$\sigma_{kl} = P_k(0)\Gamma_l. \quad (3)$$

Let $\{|k^i(s)\rangle\}_{i=1}^{n_k}$ and $\{|l^j(s)\rangle\}_{j=1}^{n_l}$ be orthonormal bases for subspaces $\mathcal{H}_k(s)$ and $\mathcal{H}_l(s)$, respectively, in terms of which

$$\begin{aligned} \sigma_{kl} &= \sum_{ij} [(\mathcal{F}_0^k | \mathcal{F}_1^l) \mathbf{P} e^{\int_0^1 \mathbf{A}_l(s) ds}]_{ij} |k^i(0)\rangle \langle l^j(0)| \\ &= \sum_{ij} [\boldsymbol{\sigma}^{kl}]_{ij} |k^i(0)\rangle \langle l^j(0)|. \end{aligned} \quad (4)$$

*Electronic address: david.kult@kvac.uu.se

†Electronic address: J.Aberg@damtp.cam.ac.uk

‡Electronic address: eriks@kvac.uu.se

Here, $(\mathcal{F}_0^k|\mathcal{F}_1^l)$ is a $n_k \times n_l$ matrix with components $(\mathcal{F}_0^k|\mathcal{F}_1^l)_{ij} = \langle k^i(0)|l^j(1)\rangle$ and $[\mathbf{A}_l(s)]_{ij} = \langle \partial_s l^i(s)|l^j(s)\rangle$ is the Wilczek-Zee connection along $\mathcal{C}_l$ in $\mathcal{G}(N; n_l)$.

The unitary part $\Phi[\sigma^{ll}][31]$ is the holonomy in ref. [24] associated with the (open) path $\mathcal{C}_l$. It seems natural to ask whether we can interpret the matrices σ^{kl} , $k \neq l$, in a similar fashion. To answer this, we need to see how these matrices behave under a gauge transformation, i.e., a change of frames $|l^i(s)\rangle \rightarrow |\tilde{l}^i(s)\rangle = \sum_j |l^j(s)\rangle [U_l(s)]_{ji}$, where $\{U_l(s)\}_{l=1}^\eta$ are unitary matrices. Under such a transformation the matrices $\mathbf{P}e^{\int_0^1 \mathbf{A}_l(s)ds}$ and $(\mathcal{F}_0^k|\mathcal{F}_1^l)$ undergo the following changes

$$\begin{aligned} \mathbf{P}e^{\int_0^1 \mathbf{A}_l(s)ds} &\rightarrow U_l^\dagger(1) \mathbf{P}e^{\int_0^1 \mathbf{A}_l(s)ds} U_l(0), \\ (\mathcal{F}_0^k|\mathcal{F}_1^l) &\rightarrow U_k^\dagger(0)(\mathcal{F}_0^k|\mathcal{F}_1^l)U_l(1). \end{aligned} \quad (5)$$

Consequently, σ^{kl} transforms as

$$\sigma^{kl} \rightarrow U_k^\dagger(0) \sigma^{kl} U_l(0), \quad (6)$$

i.e., noncovariantly unless $k = l$. Thus, the matrices σ^{kl} , $k \neq l$, fail to reflect the geometry of the paths $\mathcal{C}_k$ and $\mathcal{C}_l$. However, the specific behavior of σ^{kl} under gauge transformations suggests that we consider the operator

$$\begin{aligned} \gamma_{l_1 \dots l_\kappa} &= \sigma_{l_1 l_\kappa} \sigma_{l_\kappa l_{\kappa-1}} \dots \sigma_{l_3 l_2} \sigma_{l_2 l_1} \\ &= \sum_{ij} [\gamma^{l_1 \dots l_\kappa}]_{ij} |l_1^i(0)\rangle \langle l_1^j(0)|, \end{aligned} \quad (7)$$

where $\gamma^{l_1 \dots l_\kappa}$ is the matrix

$$\gamma^{l_1 \dots l_\kappa} = \sigma^{l_1 l_\kappa} \sigma^{l_\kappa l_{\kappa-1}} \dots \sigma^{l_3 l_2} \sigma^{l_2 l_1}. \quad (8)$$

We can use these operators and matrices to define gauge covariant quantities, since $\Phi[\gamma^{l_1 \dots l_\kappa}] \rightarrow U_{l_1}^\dagger(0) \Phi[\gamma^{l_1 \dots l_\kappa}] U_{l_1}(0)$ from eq. (6). Thus, we propose to take

$$U_g^{(\kappa)}[\mathcal{C}_{l_1}, \dots, \mathcal{C}_{l_\kappa}] = \Phi[\gamma^{l_1 \dots l_\kappa}] \quad (9)$$

as the gauge covariant non-Abelian holonomies of order κ , and thus generalizing the approach of ref. [6] to the non-Abelian case. We extend the range of κ by defining $U_g^{(1)}[\mathcal{C}_l] = \Phi[\sigma^{ll}]$, i.e., the first order ($\kappa = 1$) holonomies are taken to be the open path holonomies in refs. [23, 24].

Note that the definition in eq. (9) allows any sequence $(l_1, \dots, l_\kappa)$. This includes cases like, e.g., γ^{111} , which cannot be regarded as an “off-diagonal” object. Hence, eq. (9) can be regarded as a general definition of holonomies of degree κ , both diagonal and off-diagonal. To define genuinely off-diagonal holonomies we obtain a reasonable subclass if we require that $(l_1, \dots, l_\kappa)$ contains each number at most once. We let $\mathbb{I}_\kappa^\eta$ denote all vectors $(l_1, \dots, l_\kappa)$ with $l_j \in \{1, \dots, \eta\}$, such that none of the numbers occurs twice, e.g., $(2, 5, 3) \in \mathbb{I}_3^6$ but $(6, 4, 6, 2) \notin \mathbb{I}_4^6$. We refer to the set of holonomies $U_g^{(\kappa)}[\mathcal{C}_{l_1}, \dots, \mathcal{C}_{l_\kappa}]$ with $(l_1, \dots, l_\kappa) \in \mathbb{I}_\kappa^\eta$ with $2 \leq \kappa \leq \eta$, as “strictly off-diagonal holonomies”.

For a cyclic evolution, characterized by $\mathcal{H}_l(1) = \mathcal{H}_l(0)$, $l = 1, \dots, \eta$, the standard holonomies $U_g^{(1)}[\mathcal{C}_l]$ are fully defined. On the other hand, in this case we have $\gamma^{l_1 \dots l_\kappa} = 0$, $(l_1, \dots, l_\kappa) \in \mathbb{I}_\kappa^\eta$, $\kappa \geq 2$, which implies that all strictly off-diagonal holonomies are undefined for cyclic evolution. Thus, just as in the Abelian case [6], the standard holonomies contain all nontrivial information about $\mathcal{C}_1, \dots, \mathcal{C}_\eta$ when these are loops.

In the case where $n_l = 1$, $l = 1, \dots, \eta$, the matrices $(\mathcal{F}_0^k|\mathcal{F}_1^l)$ and $\mathbf{P}e^{\int_0^1 \mathbf{A}_l(s)ds}$ reduce to the complex numbers $\langle k(0)|l(1)\rangle$ and $e^{-\int_0^1 \langle l(s)|\partial_s l(s)\rangle ds}$, respectively. This leads to the off-diagonal geometric phase factors

$$\begin{aligned} U_g^{(\kappa)}[\mathcal{C}_{l_1}, \dots, \mathcal{C}_{l_\kappa}] &= \Phi[\langle l_1(0)|l_\kappa(1)\rangle \\ &\times e^{-\int_0^1 \langle l_\kappa(s)|\partial_s l_\kappa(s)\rangle ds} \dots \langle l_2(0)|l_1(1)\rangle \\ &\times e^{-\int_0^1 \langle l_1(s)|\partial_s l_1(s)\rangle ds}], \end{aligned} \quad (10)$$

which coincide with $\Phi[\gamma_{l_1 l_\kappa l_{\kappa-1} \dots l_2}^{(\kappa)\Gamma}]$ in ref. [6].

Manini and Pistolesi [6] suggest an interpretation of their off-diagonal geometric phases in terms of Berry phases for single closed paths. In the second order case, these paths consist of the segments $\mathcal{C}_k$, $\mathcal{C}_l$, G_{kl} , and G_{lk} , where G_{kl} geodesically connects the final point of $\mathcal{C}_k$ with the starting point of $\mathcal{C}_l$, and vice versa for G_{lk} (see fig. 1 of ref. [6]). In the general non-Abelian case, however, this interpretation is difficult to maintain. Apart from the special case when $n_1 = n_2 = \dots = n_\kappa$, it is not possible to join the curves $\mathcal{C}_{l_1}, \dots, \mathcal{C}_{l_\kappa}$, due to the mismatch of dimensions, and thus the closure using geodesics is not applicable. This observation shows that the off-diagonal holonomies in general cannot be interpreted as standard Wilczek-Zee quantum holonomies for closed paths [14], and that the off-diagonal holonomies therefore are genuinely new concepts associated with the evolution of quantum systems. Another consequence of the fact that we can have different n_l is that the rank of $\gamma^{l_1 \dots l_\kappa}$ cannot be larger than the smallest n_l (see 2.17.8 of ref. [29]), and thus may be less than n_{l_1} .

The non-Abelian character of the off-diagonal holonomies $U^{(\kappa)}[\mathcal{C}_1, \dots, \mathcal{C}_\kappa]$ implies that they are not invariant under cyclic permutations of the indexes $(l_1, \dots, l_\kappa)$. It may even be the case that two off-diagonal holonomies that differ only by a cyclic permutation have different rank, since the smallest n_l only provides an upper bound for the rank of $\gamma^{l_1 \dots l_\kappa}$. Furthermore, as is exemplified below it is also possible that $\gamma^{l_1 \dots l_\kappa}$ may have path-dependent nodal points if $\kappa \geq 2$.

III. WHERE DID THE PHASE INFORMATION GO?

As noted above, all strictly off-diagonal holonomies are undefined for cyclic evolutions, in analogy with the standard cyclic off-diagonal geometric phases. Conversely, the noncyclic geometric phases are undefined when the

states at the initial and final points of the curves are orthogonal, i.e., there exist nodal points where these phases are undefined. The hope to recover this lost phase information appears to have been one of the primary reasons for Manini and Pistoiesi to introduce off-diagonal geometric phases. However, the issue concerning the possible recovery of phase information was never explicitly investigated in ref. [6]. In the following we elucidate some aspects of this question in the case of non-Abelian off-diagonal holonomies.

Let us first analyze what happens if the rank of some of the overlap matrices $(\mathcal{F}_0^l | \mathcal{F}_1^l)$ is greater than zero but less than their subspace dimension n_l . This is a situation where the corresponding holonomies become partial [24]; a phenomenon that has no counterpart in the Abelian case. To aid us in this analysis we introduce the unitary $N \times N$ matrix

$$\mathbf{S}_{\text{tot}} = \begin{pmatrix} \sigma^{11} & \dots & \sigma^{1\eta} \\ \vdots & \ddots & \vdots \\ \sigma^{\eta 1} & \dots & \sigma^{\eta\eta} \end{pmatrix}. \quad (11)$$

It follows from unitarity that $R(\mathbf{S}_{\text{tot}}) = N$, where $R(\mathbf{X})$ denotes the rank of matrix $\mathbf{X}$. Furthermore, for every $l = 1, \dots, \eta$, it holds that $\sum_k \sigma^{kl\dagger} \sigma^{kl} = \sum_k \sigma^{lk} \sigma^{lk\dagger} = \mathbf{1}_{n_l \times n_l}$, where $\mathbf{1}_{n_l \times n_l}$ denotes the $n_l \times n_l$ identity matrix. This entails that (see 2.17.2 and 2.17.5 of ref. [29]) $\sum_k R(\sigma^{kl}) \geq n_l$ and $\sum_k R(\sigma^{lk}) \geq n_l$. So, if $R(\sigma^{ll}) = n_l - n$, then $\sum_{k \neq l} R(\sigma^{kl}) \geq n$ and $\sum_{k \neq l} R(\sigma^{lk}) \geq n$. In other words, when the overlap matrix $(\mathcal{F}_0^l | \mathcal{F}_1^l)$ decreases by n in rank, the lower bound for the sum of the ranks of the matrices σ^{kl} increases by the same amount. Thus, the ‘‘holonomy information’’ that is lost when the holonomy of the curve $\mathcal{C}_l$ becomes partial is transferred to the matrices σ^{kl} .

A perhaps more significant question is whether the rank of the matrices $\gamma^{l_1 \dots l_\kappa}$ depends on the rank of the overlap matrix $(\mathcal{F}_0^l | \mathcal{F}_1^l)$ in a manner similar to what was discussed above for the matrices σ^{kl} . One can demonstrate with a simple counterexample that no such relation exists. Assume $\eta = 3$ and $n_1 = n_2 = n_3 = 2$. Furthermore, assume that σ^{13} , σ^{21} , and σ^{32} are the 2×2 zero matrix, and

$$\sigma^{11\dagger} = \sigma^{22\dagger} = \sigma^{33\dagger} = \sigma^{12} = \sigma^{23} = \sigma^{31} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

One may verify that the corresponding matrix $\mathbf{S}_{\text{tot}}$ is unitary. In this example $\gamma^{kl} = 0$, $\forall k, l$, and $\gamma^{klm} = 0$, $\forall k, l, m$. Hence, although none of the overlap matrices $(\mathcal{F}_0^l | \mathcal{F}_1^l)$ are of full rank, all strictly off-diagonal holonomies vanish, even those of higher order.

Finally, we ask what happens if the matrices σ^{ll} are zero for all $l = 1, \dots, \eta$. We prove by *reductio ad absurdum* that at least one of the strictly off-diagonal holonomies must have nonzero rank. Assume $\sigma^{ll} = 0$, for $l = 1, \dots, \eta$, and $\gamma^{l_1 \dots l_\kappa} = 0$, for $(l_1, \dots, l_\kappa) \in \mathbb{I}_\kappa^\eta$, $\kappa \geq 2$. Consider an arbitrary string $(l_1, \dots, l_\nu)$ consisting of the integers 1 to η , and with $\nu \geq 2$. If

$(l_1, \dots, l_\nu) \in \mathbb{I}_\nu^\eta$ ($\nu \leq \eta$), then by assumption $\gamma^{l_1 \dots l_\nu} = 0$. If $(l_1, \dots, l_\nu) \notin \mathbb{I}_\nu^\eta$, then take one of the smallest subsequences $(l_\alpha, l_{\alpha+1}, \dots, l_{\beta-1}, l_\beta)$ that begins and ends with the same number (i.e., $l_\beta = l_\alpha$). In this subsequence there is no other repetitions (otherwise there exists a smaller subsequence). It follows that $\gamma^{l_1 \dots l_\nu} = \sigma^{l_1 l_\nu} \dots \sigma^{l_{\beta+1} l_\beta} \gamma^{l_\alpha \dots l_{\beta-1}} \sigma^{l_\alpha l_{\alpha-1}} \dots \sigma^{l_2 l_1}$. Moreover, $(l_\alpha, \dots, l_{\beta-1}) \in \mathbb{I}_{\beta-\alpha+1}^\eta$. If $\beta - \alpha = 1$, then $\gamma^{l_\alpha \dots l_{\beta-1}} = \sigma^{l_\alpha l_\alpha} = 0$, otherwise $\gamma^{l_\alpha \dots l_{\beta-1}} = 0$ by assumption. We proceed by noting that $\text{Tr}(\mathbf{S}_{\text{tot}}^\nu) = \sum_{l_1 \dots l_\nu} \text{Tr} \gamma^{l_1 \dots l_\nu} = 0$, for all $\nu = 1, 2, \dots$, as a consequence of our assumptions. However, this cannot be the case since $\mathbf{S}_{\text{tot}}$ is a unitary matrix. Therefore, our assumptions must be wrong, and at least one of the strictly off-diagonal holonomies must have nonzero rank. By the above findings we can conclude that the holonomy information lost in the the nodal points of the non-Abelian noncyclic holonomies indeed can be retained in some sense, but that the structure is much more intricate and rich than in the Abelian case, due to the existence of partial holonomies in the non-Abelian setting.

IV. EXAMPLE

We illustrate the off-diagonal holonomies by an example in the adiabatic context. We let the system evolve under the action of a slowly varying Hamiltonian. Let us consider the tripod system [18, 30] modeled by the parameter-dependent four-state Hamiltonian $H(s) = \omega|e\rangle(\sin\theta(s)\cos\varphi(s)\langle 0| + \sin\theta(s)\sin\varphi(s)\langle 1| + \cos\theta(s)\langle a|) + \text{h.c.}$, exhibiting two nondegenerate ‘bright’ states $|F_s^\pm\rangle = \{|B^\pm(s)\rangle\}$ with energy $\pm\omega$ and a doubly degenerate ‘dark’ zero energy eigenspace $|F_s^d\rangle = \{|D^1(s)\rangle, |D^2(s)\rangle\}$. Explicitly, we may choose

$$\begin{aligned} |B^\pm\rangle &= \frac{1}{\sqrt{2}}(|e\rangle \pm \sin\theta\cos\varphi|0\rangle \pm \sin\theta\sin\varphi|1\rangle \\ &\quad \pm \cos\theta|a\rangle), \\ |D_1\rangle &= \cos\theta\cos\varphi|0\rangle + \cos\theta\sin\varphi|1\rangle - \sin\theta|a\rangle, \\ |D_2\rangle &= -\sin\varphi|0\rangle + \cos\varphi|1\rangle. \end{aligned} \quad (12)$$

Consider paths $(0, 0) \rightarrow (\theta_1, \varphi_1)$ in parameter space (θ, φ) . For each such path the energy eigenstates define paths $\mathcal{C}_\pm$ in $\mathcal{G}(4; 1)$ and $\mathcal{C}_d$ in $\mathcal{G}(4; 2)$. We obtain the geometric phase factors $\mathbf{U}^{(1)}[\mathcal{C}_\pm] = 1$ for $\theta_1 \neq \pi$ and $\mathbf{U}^{(2)}[\mathcal{C}_\pm, \mathcal{C}_\mp] = 1$ for $\theta_1 \neq 0$. $\mathbf{U}^{(1)}[\mathcal{C}_\pm]$ are undefined at $\theta_1 = \pi$ and similarly $\mathbf{U}^{(2)}[\mathcal{C}_\pm, \mathcal{C}_\mp]$ at $\theta_1 = 0$. In ref. [24] it was shown that $\mathbf{U}^{(1)}[\mathcal{C}_d]$ is fully defined, except when the path ends at $\theta_1 = \pi/2$, where the holonomy becomes partial. The strictly off-diagonal holonomies ($\kappa = 2, 3$) involving the dark subspace are undefined when $\sin\theta_1 = 0$.




FIG. 1: Interferometric approach to obtain the holonomy $U_g^{(\kappa)}[C_{l_1}, \dots, C_{l_\kappa}]$. The horizontal and vertical paths of the Mach-Zehnder interferometer correspond to the states $|0\rangle$ and $|1\rangle$, respectively. The particle have an internal degree of freedom, on which the unitary operation U is applied. U can be generated either by adiabatic or nonadiabatic evolution, or approximated by filtering measurements. The particle enters the interferometer in path 0 and in the internal state $P_{l_1}(0)/n_{l_1}$. In path 1 a variable unitary operator V such that $[V, P_l(0)] = 0$, $\forall l$, is applied, while in path 0 we alternately apply U and a filtering corresponding to the projector $|0\rangle\langle 0| \otimes P_{l_k}(0) + |1\rangle\langle 1| \otimes \hat{1}$, for $k = 2, \dots, \kappa$. A final U is applied, followed by the filtering $|0\rangle\langle 0| \otimes P_{l_1}(0) + |1\rangle\langle 1| \otimes \hat{1}$. The probability p to find the particle in path 0 is measured after a second beam splitter. The unitary operator V is varied so as to maximize p .

For $\sin \theta_1 \neq 0$, let $Z = \int_0^1 \cos[\theta(s)] \dot{\varphi}(s) ds$ and we obtain

$$\begin{aligned} U^{(2)}[C_\pm, C_d] &= -U^{(3)}[C_\pm, C_\mp, C_d] \\ &= -U^{(3)}[C_\pm, C_d, C_\mp] = -\frac{\cos(\varphi_1 - Z)}{|\cos(\varphi_1 - Z)|}, \\ U^{(2)}[C_d, C_\pm] &= -U^{(3)}[C_d, C_\pm, C_\mp] \\ &= -\begin{pmatrix} \cos Z \cos \varphi_1 & \sin Z \cos \varphi_1 \\ \cos Z \sin \varphi_1 & \sin Z \sin \varphi_1 \end{pmatrix}. \end{aligned} \quad (13)$$

While $U^{(2)}[C_d, C_\pm]$ and $U^{(3)}[C_d, C_\pm, C_\mp]$ are nonzero partial isometries, there are path-dependent nodal points of $U^{(2)}[C_\pm, C_d]$, $U^{(3)}[C_\pm, C_\mp, C_d]$, and $U^{(3)}[C_\pm, C_d, C_\mp]$, namely where $\cos(\varphi_1 - Z) = 0$.

V. PHYSICAL REALIZATIONS

Let us now examine some possible physical realizations of $U_g^{(\kappa)}[C_{l_1}, \dots, C_{l_\kappa}]$. Consider the Mach-Zehnder interferometer in fig. 1 with the two path states represented by $|0\rangle$ and $|1\rangle$. We let the internal state of the particle (e.g., spin) be represented by the Hilbert space $\mathcal{H} = \mathcal{H}_1(s) \oplus \dots \oplus \mathcal{H}_\eta(s)$, $s \in [0, 1]$. The total system is prepared in the state $|0\rangle\langle 0| \otimes P_{l_1}(0)/n_{l_1}$. We first apply a beam-splitter, followed by the unitary operations $|0\rangle\langle 0| \otimes \hat{1} + |1\rangle\langle 1| \otimes V$ and $|0\rangle\langle 0| \otimes U + |1\rangle\langle 1| \otimes \hat{1}$, where V is a variable unitary operator assumed to be chosen such that $[V, P_l(0)] = 0$ for all l . Next, we perform a filtering corresponding to the projection operator

$|0\rangle\langle 0| \otimes P_{l_2}(0) + |1\rangle\langle 1| \otimes \hat{1}$, i.e., the particle is “removed” if it is found in path 0 with its internal state outside subspace $\mathcal{H}_{l_2}$. Thereafter, we again apply the operator $|0\rangle\langle 0| \otimes U + |1\rangle\langle 1| \otimes \hat{1}$, and the filtering $|0\rangle\langle 0| \otimes P_{l_3}(0) + |1\rangle\langle 1| \otimes \hat{1}$. This procedure is repeated until we have applied the operator $|0\rangle\langle 0| \otimes U + |1\rangle\langle 1| \otimes \hat{1}$, κ times. After this, we apply a final filtering $|0\rangle\langle 0| \otimes P_{l_1}(0) + |1\rangle\langle 1| \otimes \hat{1}$, and recombine the two paths with a beam splitter. We finally measure the probability p to find the particle in path 0.

First, we assume that the unitary operator U acting on $\mathcal{H}$ is caused by an adiabatic evolution of a time-dependent Hamiltonian with eigenspaces $\{\mathcal{H}_l(s)\}_{l=1}^\eta$. This allows us to write $U = \sum_l e^{i\phi_l} \Gamma_l$, where ϕ_l is the dynamical phase $\phi_l = \int_0^1 E_l(s) ds$, and $E_l(s)$ the eigenvalue corresponding to eigenspace $\mathcal{H}_l(s)$ of the Hamiltonian. The corresponding detection probability becomes

$$\begin{aligned} p &= \frac{1}{4} + \frac{1}{4} \frac{1}{n_{l_1}} \text{Tr}(\gamma_{l_1 \dots l_\kappa} \gamma_{l_1 \dots l_\kappa}^\dagger) \\ &+ \frac{1}{2} \frac{1}{n_{l_1}} \text{Re}[e^{i \sum_{k=1}^\kappa \phi_k} \text{Tr}(\gamma^{l_1 \dots l_\kappa} \mathbf{V}^\dagger)], \end{aligned} \quad (14)$$

where $\mathbf{V}_{ij} = \langle l_1^i(0) | V | l_1^j(0) \rangle$. Note that $\mathbf{V}$ is a unitary matrix since $[V, P_{l_1}(0)] = 0$. By varying V we obtain the maximal detection probability when $\mathbf{V} = e^{i \sum_{k=1}^\kappa \phi_k} U_g^{(\kappa)}[C_{l_1}, \dots, C_{l_\kappa}]$. Hence, up to the dynamical phases we have found the holonomy.

In the adiabatic setting there is in the general case no easy way to eliminate the dynamical phases. To avoid these problems we consider two alternative approaches to generate the unitary operator U . One alternative is to base the evolution entirely on filtering, where we approximate the evolution in the spirit of ref. [22]. We begin with the same initial state, beam-splitter, and variable unitary V , as in the previous case. Next we apply a sequence of filterings $|0\rangle\langle 0| \otimes P_{l_1}(s_j) + |1\rangle\langle 1| \otimes \hat{1}$, where s_j form a discretization of the interval $[0, 1]$. For the next step we apply the sequence of filterings $|0\rangle\langle 0| \otimes P_{l_2}(s_j) + |1\rangle\langle 1| \otimes \hat{1}$, and we continue up l_κ . We finally apply $|0\rangle\langle 0| \otimes P_{l_1}(0) + |1\rangle\langle 1| \otimes \hat{1}$, followed by a beam splitter, and measure the probability to find the particle in path 0. One can show that the probability is

$$\begin{aligned} p &= \frac{1}{4} + \frac{1}{4} \frac{1}{n_{l_1}} \text{Tr}(\gamma_{l_1 \dots l_\kappa} \gamma_{l_1 \dots l_\kappa}^\dagger) \\ &+ \frac{1}{2} \frac{1}{n_{l_1}} \text{Re}[\text{Tr}(\gamma^{l_1 \dots l_\kappa} \mathbf{V}^\dagger)]. \end{aligned} \quad (15)$$

Hence, as in eq. (14), apart from the absence of dynamical phases.

The second alternative that allows us to avoid the problem with dynamical phases is to use a nonadiabatic approach. Assume that the evolution is driven by the time-dependent Hamiltonian $H(s)$, where now s is the time-parameter. We let the subspaces $\mathcal{H}_l(s)$ be evolving under $H(s)$ according to the Schrödinger equation. In contrast to the adiabatic approach the subspaces $\mathcal{H}_l(s)$

are in the general case not eigenspaces of $H(s)$. We furthermore let $\{|l^j(s)\rangle\}_{j=1}^{n_l}$ be smoothly parameterized orthonormal bases of the subspaces $\mathcal{H}_l(s)$. We wish to find the unitary matrices $\mathbf{U}_l(s)$ such that the vectors

$$|\chi_l^k(s)\rangle = \sum_j |l^j(s)\rangle [\mathbf{U}_l(s)]_{jk} \quad (16)$$

satisfy the Schrödinger equation $i\partial_s |\chi_l^k(s)\rangle = H(s) |\chi_l^k(s)\rangle$ ($\hbar = 1$) with initial conditions $|\chi_l^k(0)\rangle = |l^k(0)\rangle$. If we substitute eq. (16) into the Schrödinger equation we find that $\mathbf{U}_l(s)$ has to satisfy $i\partial_s \mathbf{U}_l(s) = i\mathbf{A}_l(s)\mathbf{U}_l(s) + \mathbf{K}_l(s)\mathbf{U}_l(s)$, where $[\mathbf{A}_l(s)]_{j'j} = \langle \partial_s l^{j'}(s) | l^j(s) \rangle$ and $[\mathbf{K}_l(s)]_{j'j} = \langle l^{j'}(s) | H(s) | l^j(s) \rangle$ contain the geometrical and dynamical contributions, respectively, as was discussed in ref. [21]. In order to get rid of the dynamical contribution without affecting the evolution of the subspaces we introduce the modified time-dependent Hamiltonian

$$\overline{H}(s) = H(s) - \sum_{l=1}^{\eta} P_l(s) H(s) P_l(s). \quad (17)$$

The evolution of the subspaces $\mathcal{H}_l(s)$ are not affected by this modification since $[\overline{H}(s), P_l(s)] = [H(s), P_l(s)]$, $\forall l$. We now wish to find the unitary matrices $\overline{\mathbf{U}}_l(s)$ such that the vectors $|\overline{\chi}_l^k(s)\rangle = \sum_j |l^j(s)\rangle [\overline{\mathbf{U}}_l(s)]_{jk}$ satisfy the modified Schrödinger equation $i\partial_s |\overline{\chi}_l^k(s)\rangle = \overline{H}(s) |\overline{\chi}_l^k(s)\rangle = [H(s) - P_l(s)H(s)P_l(s)] |\overline{\chi}_l^k(s)\rangle$ with initial conditions $|\overline{\chi}_l^k(0)\rangle = |l^k(0)\rangle$. In this case we obtain $\partial_s \overline{\mathbf{U}}_l(s) = i\mathbf{A}_l(s)\overline{\mathbf{U}}_l(s)$. Hence, the solution $\overline{\mathbf{U}}_l(s) = \mathbf{P}e^{\int_0^s \mathbf{A}_l(s') ds'}$ only depends on the geometric contribution[32]. The time-dependent Hamiltonian $\overline{H}(s)$ generates a unitary mapping from the initial state to the state at time $s = 1$ that is given by $\overline{U}(1) = \sum_l \Gamma_l$, where

$$\begin{aligned} \Gamma_l &= \sum_k |\overline{\chi}_l^k(1)\rangle \langle \overline{\chi}_l^k(0)| \\ &= \sum_{jk} [\mathbf{P}e^{\int_0^1 \mathbf{A}_l(s) ds}]_{jk} |l^j(1)\rangle \langle l^k(0)|. \end{aligned} \quad (18)$$

This means that if we let $U = \overline{U}(1)$ in the alternating procedure described above (see fig. 1), then the probability to detect the particle in path 0 becomes as in eq. (15), and is maximized when $\mathbf{V} = \mathbf{U}_g^{(\kappa)}[\mathcal{C}_{l_1}, \dots, \mathcal{C}_{l_\kappa}]$.

VI. CONCLUSION

Noncyclic evolution of quantum systems may lead to well-defined off-diagonal holonomies that involve more than one subspace of Hilbert space. These holonomies reduce to the off-diagonal geometric phases in ref. [6] for one-dimensional subspaces. The off-diagonal holonomies are undefined for cyclic evolution but must contain members of nonzero rank when all the standard holonomies are undefined. While the nodal point structure of the holonomy for an open continuous path [24] can only depend on the end-points of the path, this structure can be path-dependent in the off-diagonal case. Furthermore, we have put forward physical realizations of the off-diagonal holonomies in the context of adiabatic evolution and interferometry that may open up the possibility to test these quantities experimentally.

Acknowledgments

J.Å. thanks the Swedish Research Council for financial support and the Centre for Quantum Computation at DAMTP, Cambridge, for hospitality. E.S. acknowledges financial support from the Swedish Research Council.

-
- [1] SAMUEL J. and BHANDARI R., *Phys. Rev. Lett.*, **60** (1988) 2339.
 - [2] BHANDARI R., *Phys. Lett. A*, **157** (1991) 221.
 - [3] BHANDARI R., *Phys. Lett. A*, **171** (1992) 262.
 - [4] BHANDARI R., *Phys. Lett. A*, **171** (1992) 267.
 - [5] BHANDARI R., *Phys. Lett. A*, **180** (1993) 15.
 - [6] MANINI N. and PISTOLESI F., *Phys. Rev. Lett.*, **85** (2000) 3067.
 - [7] MUKUNDA N., ARVIND, CHATURVEDI S. and SIMON R., *Phys. Rev. A*, **65** (2002) 012102.
 - [8] HASEGAWA Y., LOIDL R., BARON M., BADUREK G. and RAUCH H., *Phys. Rev. Lett.*, **87** (2001) 070401.
 - [9] HASEGAWA Y., LOIDL R., BADUREK G., BARON M., MANINI N., PISTOLESI F. and RAUCH H., *Phys. Rev. A*, **65** (2002) 052111.
 - [10] FILIPP S. and SJÖQVIST E., *Phys. Rev. Lett.*, **90** (2003) 050403 (2003).
 - [11] FILIPP S. and SJÖQVIST E., *Phys. Rev. A*, **68** (2003) 042112.
 - [12] TONG D. M., SJÖQVIST E., FILIPP S., KWEK L. C. and OH C. H., *Phys. Rev. A*, **71** (2005) 032106.
 - [13] FILIPP S. and SJÖQVIST E., *Phys. Lett. A*, **342** (2005) 205.
 - [14] WILCZEK F. and ZEE A., *Phys. Rev. Lett.*, **52** (1984) 2111.
 - [15] ZANARDI P. and RASETTI M., *Phys. Lett. A*, **264** (1999) 94.
 - [16] PACHOS J., ZANARDI P. and RASETTI M., *Phys. Rev. A*, **61** (2000) 010305(R).
 - [17] PACHOS J. and CHOUNTASIS S., *Phys. Rev. A*, **62**

- (2000) 052318.
- [18] DUAN L. -M., CIRAC J. I. and ZOLLER P., *Science*, **292** (2001) 1695.
 - [19] RECATI A., CALARCO T., ZANARDI P., CIRAC J. I. and ZOLLER P., *Phys. Rev. A*, **66** (2002) 032309.
 - [20] FAORO L., SIEWERT J. and FAZIO R., *Phys. Rev. Lett.*, **90** (2003) 028301.
 - [21] ANANDAN J., *Phys. Lett. A*, **133** (1988) 171.
 - [22] SJÖQVIST E., KULT D. and ÅBERG J., *Phys. Rev. A*, **74** (2006) 062101.
 - [23] MOSTAFAZADEH A., *J. Phys. A*, **32** (1999) 8157.
 - [24] KULT D., ÅBERG J. and SJÖQVIST E., *Phys. Rev. A*, **74** (2006) 022106.
 - [25] GREUB W., HALPERIN S. and VANSTONE R., *Connections, Curvature and Cohomology* Vol. II (Academic Press, New York) 1973.
 - [26] MOORE E. H., *Bull. Am. Math. Soc.*, **26** (1920) 394.
 - [27] PENROSE R., *Proc. Cambridge Phil. Soc.*, **51** (1955) 406.
 - [28] LANCASTER P. and TISMENETSKY M., *The Theory of Matrices* (Academic Press, San Diego) 1985.
 - [29] MARCUS M. and MINC H., *A Survey of Matrix Theory and Matrix Inequalities* (Allyn and Bacon, Inc, Boston) 1964.
 - [30] UNANYAN R. G., SHORE B. W. and BERGMANN K., *Phys. Rev. A*, **59** (1999) 2910.
 - [31] For a matrix $\mathbf{Z}$, $\Phi[\mathbf{Z}] \equiv (\sqrt{\mathbf{Z}\mathbf{Z}^\dagger})^\ominus \mathbf{Z}$, $\ominus$ being the Moore-Penrose pseudoinverse [26, 27, 28] obtained by inverting all nonzero eigenvalues of $\sqrt{\mathbf{Z}\mathbf{Z}^\dagger}$. If $\det \sqrt{\mathbf{Z}\mathbf{Z}^\dagger} \neq 0$ then the MP pseudoinverse of $\sqrt{\mathbf{Z}\mathbf{Z}^\dagger}$ coincides with the inverse.
 - [32] One may note, though, that $\overline{\mathbf{U}}_l(s)$ is not gauge covariant and can therefore not be considered a geometric quantity, see ref. [24].